



# Radiocarbon Dating of Carbonised Plant Macrofossils

Woodcutts, Iwerne, Rotherley, Durrington Walls, Cuckoo Stone, Coneybury Henge, Lockington, Thanet Earth and Hunsbury Hillfort

Peter Marshall, Ruth Pelling, Christopher Bronk Ramsey, Elaine Dunbar, Irka Hajdas, Sanne Palstra and Paula Reimer



# Radiocarbon Dating of Carbonised Plant Macrofossils

Woodcutts, Iwerne, Rotherley, Durrington Walls,  
Cuckoo Stone, Coneybury Henge, Lockington, Thanet  
Earth and Hunsbury Hillfort

Peter Marshall, Ruth Pelling, Christopher Bronk Ramsey, Elaine Dunbar, Irka Hajdas,  
Sanne Palstra and Paula Reimer

2023

NGR: ST 963 181, ST 856 137, ST 9491 1950, SU 14955 43598, SU 146 433, SU  
134 416, SK 4690 2910, TR 2878 6680, SP 73792 58361

Print: ISSN 2398-3841

Online: ISSN 2059-4453

The Research Report Series incorporates reports by Historic England's expert teams, their predecessors and other researchers. Many Research Reports are interim, to make available the results of specialist investigations in advance of full publication. Although subject to internal quality assurance, they are not usually refereed externally and their conclusions may sometimes have to be modified in the light of information not available at the time of the investigation. Where no final project report is available, readers should consult the author before citing these reports.

For more information email [Res.reports@HistoricEngland.org.uk](mailto:Res.reports@HistoricEngland.org.uk) or write to:

Historic England, Fort Cumberland, Fort Cumberland Road, Eastney,  
Portsmouth PO4 9LD

Opinions expressed in Research Reports are those of the author(s) and are not necessarily those of Historic England.

## Summary

This document is a technical archive report on the radiocarbon dating of carbonised plant macrofossils in support of research undertaken by the Historic England Environmental Studies branch. It includes full details of 32 radiocarbon measurements obtained from Woodcutts, Iwerne, Rotherley, Durrington Walls, Cuckoo Stone, Coneybury Henge, Lockington, Thanet Earth and Hunsbury Hillfort.

## Contributors

Peter Marshall, Ruth Pelling, Christopher Bronk Ramsey, Elaine Dunbar, Irka Hajdas, Sanne Palstra and Paula Reimer.

## Acknowledgements

We would like to thank Wendy Carruthers, Angela Monkton and Ellen Simmons, for providing material for dating. The front cover photo shows a box of charred grain from Rotherley pit 43 without its lid from the Pitt Rivers collection held at Salisbury Museum and was taken at Fort Cumberland by Steven Baker (Historic England) on the 23 January 2017.

## Archive location

Historic England, The Engine House, Fire Fly Avenue, Swindon, SN2 2EH.

## Historic environment records

Wiltshire Archaeology Service, The Wiltshire and Swindon History Centre, Cocklebury Road, Chippenham, SN15 3QN.

Northamptonshire Record Office, Wootton Hall Park, Mereway, Northampton, Northamptonshire, NN4 8BQ.

Heritage Conservation, Kent County Council, Invicta House, County Hall, Maidstone, Kent, ME14 1XX.

Historic & Natural Environment Team, Leicestershire County Council, Room 200, County Hall, Glenfield, Leicestershire, LE3 8RA.

## Date of research

2013–2023

## Contact details

Historic England. Cannon Bridge House, 25 Dowgate Hill, London, EC4R 2YA. 020 7973 3700. [customers@HistoricEngland.org.uk](mailto:customers@HistoricEngland.org.uk).

Peter Marshall, Historic England, Cannon Bridge House, 25 Dowgate Hill, London, EC4R 2YA. [peter.marshall@historicengland.org.uk](mailto:peter.marshall@historicengland.org.uk)

Ruth Pelling, Historic England, Fort Cumberland, Fort Cumberland Road, Portsmouth, Hampshire, PO4 9LD. [ruth.pelling@historicengland.org.uk](mailto:ruth.pelling@historicengland.org.uk)

Christopher Bronk Ramsey, Oxford Radiocarbon Accelerator Unit, Research Laboratory for Archaeology and the History of Art, South Parks Road, Oxford, OX1 3QY. [christopher.ramsey@rlaha.ox.ac.uk](mailto:christopher.ramsey@rlaha.ox.ac.uk)

Elaine Dunbar, Scottish Universities Environmental Research Centre, Scottish Enterprise Technology Park, Rankine Avenue, East Kilbride G75 0QF. [elaine.dunbar@glasgow.ac.uk](mailto:elaine.dunbar@glasgow.ac.uk)

Irka Hajdas, Laboratory of Ion Beam Physics, ETH Zürich, Otto-Stern-Weg 5, CH-8093 Zürich, Switzerland. [hajdas@phys.ethz.ch](mailto:hajdas@phys.ethz.ch)

Sanne Palstra, Centre for Isotope Research, University of Groningen, Nijenborgh 6, 9747 AG Groningen, The Netherlands. [s.w.l.palstra@rug.nl](mailto:s.w.l.palstra@rug.nl)

Paula Reimer, <sup>14</sup>CHRONO Centre, Queen's University Belfast, 42 Fitzwilliam Street, Belfast, BT9 6AX. [p.j.reimer@qub.ac.uk](mailto:p.j.reimer@qub.ac.uk)

# Contents

|                                                                    |    |
|--------------------------------------------------------------------|----|
| Introduction.....                                                  | 1  |
| Radiocarbon dating .....                                           | 2  |
| Laboratory methods.....                                            | 2  |
| Calibration .....                                                  | 3  |
| Results.....                                                       | 4  |
| Pitt Rivers archive.....                                           | 4  |
| Neolithic and Bronze Age cereals in the Stonehenge landscape ..... | 7  |
| Contamination: archaeophytes .....                                 | 13 |
| Regional Reviews of charred plant remains .....                    | 15 |
| Hunsbury .....                                                     | 16 |
| References .....                                                   | 18 |

# Illustrations

|                                                                                                                                                                                                                                                |    |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----|
| Figure 1: Sites with radiocarbon dated carbonised plant macrofossils .....                                                                                                                                                                     | 1  |
| Figure 2: Probability distribution of the dates on carbonised cereal grains from the Pitt Rivers archive (Woodcutts, Iwerne and Rotherley). The distributions are the result of simple radiocarbon calibration (Stuiver and Reimer 1993) ..... | 6  |
| Figure 3: Probability distribution of the dates on carbonised cereal grains from Durrington Walls. The distributions are the result of simple radiocarbon calibration (Stuiver and Reimer 1993) .....                                          | 8  |
| Figure 4: Probability distribution of the dates on carbonised cereal grains from the Cuckoo Stone. The distributions are the result of simple radiocarbon calibration (Stuiver and Reimer 1993) .....                                          | 10 |
| Figure 5: Probability distribution of the dates on carbonised cereal grains from Coneybury Henge. The distributions are the result of simple radiocarbon calibration (Stuiver and Reimer 1993) .....                                           | 12 |
| Figure 6: Probability distribution of the date on plant macrofossils from Thanet Earth. The distribution is the result of simple radiocarbon calibration (Stuiver and Reimer 1993) .....                                                       | 14 |
| Figure 7: Probability distribution of the dates from Lockington F182. The distributions are the result of simple radiocarbon calibration (Stuiver and Reimer 1993) .....                                                                       | 15 |
| Figure 8: Probability distribution of the dates on carbonised cereal grains from Hunsbury hillfort. The distributions are the result of simple radiocarbon calibration (Stuiver and Reimer 1993) .....                                         | 16 |

# Tables

|                                                                                                              |    |
|--------------------------------------------------------------------------------------------------------------|----|
| Table 1: Pitt Rivers archive: Woodcutts, Iwerne, and Rotherley – radiocarbon and stable isotope results..... | 6  |
| Table 2: Durrington Walls radiocarbon and stable isotope results .....                                       | 8  |
| Table 3: Cuckoo Stone radiocarbon and stable isotope measurements .....                                      | 11 |
| Table 4: Coneybury Henge radiocarbon and stable isotope measurements.....                                    | 12 |
| Table 5: Thanet Earth radiocarbon and stable isotope measurements.....                                       | 14 |
| Table 6: Lockington radiocarbon and stable isotope measurements .....                                        | 16 |
| Table 7: Hunsbury radiocarbon and stable isotope measurements .....                                          | 16 |

## Introduction

This document is a technical archive report on the radiocarbon dating of 32 carbonised plant macrofossils from Woodcutts, Iwerne, Rotherley, and Hunsbury Hillfort (all from historic archives), Durrington Walls, Cuckoo Stone, Coneybury henge, Lockington, and Thanet Earth (Fig. 1), in support of research undertaken by the Historic England Environmental Studies team. It is beyond the scope of this document to detail the significance or otherwise of the results, however, elements of this report may be combined with further information in the future as part of comprehensive publications.




Figure 1: Sites with radiocarbon dated carbonised plant macrofossils

# Radiocarbon dating

## Laboratory methods

Radiocarbon measurements were made on a total of 32 plant macrofossils at the Oxford Radiocarbon Accelerator Unit (ORAU), Scottish Universities Environmental Research Centre (SUERC) <sup>14</sup>CHRON centre, The Queen's University Belfast, Centre for Isotope Research, University of Groningen (CIO) and Laboratory of Ion Beam Physics, ETH Zürich.

The eight samples submitted to the ORAU were pre-treated following methods outlined in Brock et al. (2010), graphitised (Dee and Bronk Ramsey 2000) and dated by Accelerator Mass Spectrometry (AMS) (Bronk Ramsey et al. 2004).

The 10 samples dated at The Queen's University Belfast were processed using an acid wash (Reimer et al. 2015). The pre-treated and dried samples were placed in quartz tubes with a strip of silver ribbon to remove nitrates, chlorides, and CuO. The samples were then sealed under vacuum and combusted to CO<sub>2</sub> overnight at 850°C. The CO<sub>2</sub> was converted to graphite on an iron catalyst using the zinc reduction method (Vogel et al. 1984). The graphite samples were analysed with a 0.5MeV NEC pelletron compact accelerator, with the <sup>14</sup>C/<sup>12</sup>C ratios corrected for fractionation using the on-line measured <sup>13</sup>C/<sup>12</sup>C ratio and in accordance with Stuiver and Polach (1977).

The seven samples dated at SUERC were pre-treated and dated by AMS as described by Dunbar et al. (2016).

Three samples were dated at CIO. They were pre-treated using an acid-base-acid protocol (4% HCl, 1% NaOH, <1% HCl) and combusted in an elemental analyser (IsotopeCube NCS), coupled to an Isotope Ratio Mass Spectrometer (Isoprime 100) (Dee et al. 2019). The resultant CO<sub>2</sub> was graphitised by hydrogen reduction in the presence of an iron catalyst. The graphite was then pressed into aluminium cathodes and dated by AMS (Synal et al. 2007; Salehpour et al. 2016).

Four samples were dated at ETH Zürich. They were pre-treated using the acid-base-acid protocol described by Hajdas (2008), combusted and graphitised as outlined in Wacker et al. (2010a), and dated by AMS (Synal et al. 2007; Wacker et al. 2010b).

Data reduction at the CIO and ETH Zürich was undertaken as described by Wacker et al. (2010c).

All five facilities maintain continual programmes of quality assurance procedures, in addition to participation in international inter-comparison exercises (Scott et al. 2017). Details of quality assurance data and error calculation at Groningen are provided by Aerts-Bijma et al. (2021), and similar details for ETH are provided in Sookdeo et al. (2020).

## Calibration

Radiocarbon dating is based on the radioactive decay of  $^{14}\text{C}$ , which trees absorb from the atmosphere during photosynthesis and store in their growth-rings. The radiocarbon from each year is stored in a separate annual ring. Once a ring has formed, no more  $^{14}\text{C}$  is added to it, and so the proportion of  $^{14}\text{C}$  versus other carbon isotopes reduces in the ring through time as the radiocarbon decays. Radiocarbon ages, like those in Tables 1–7, measure the proportion of  $^{14}\text{C}$  in a sample and are expressed in radiocarbon years BP (before present, ‘present’ being a constant, conventional date of AD 1950).

Radiocarbon ages are not the same as calendar dates because the concentration of  $^{14}\text{C}$  in the atmosphere has fluctuated over time. A radiocarbon measurement has thus to be calibrated against an independent scale to arrive at the corresponding calendar date. That independent scale is the IntCal20 calibration curve (Reimer et al. 2020). For the period covered by the samples dated in this report, this is constructed from radiocarbon measurements on tree-ring samples dated absolutely by dendrochronology. The probability distributions of the calibrated radiocarbon dates derived from the probability method (Stuiver and Reimer 1993) using the program OxCal v4.4 (<http://c14.arch.ox.ac.uk/oxcal.html>; Bronk Ramsey 1995; 2001; 2009). are shown in Figures 2–8. The date ranges given in Tables 1–7 have been calculated using the probability method (Stuiver and Reimer 1993), and are quoted in the form recommended by Mook (1986); rounded outwards to 10 years if the error is >25 years and five years if the error is <25 years.

# Results

## Pitt Rivers archive

Known as the ‘father of modern scientific archaeology’, General Pitt Rivers spent the last twenty years of his life (from 1880–1900) excavating sites on his estate on Cranborne Chase. The collection of artefacts found during this period was given to the Salisbury and Wiltshire Museum in 1975 by HM Treasury in lieu of death duty (<https://collections.salisburymuseum.org.uk/> Pitt Rivers Collection). Among the collection was a series of small boxes containing charred cereal remains, still retaining their original labels, as well as a selection of charcoal, some of which have been on display in Salisbury Museum for some years, although none of the material had previously been examined using modern criteria. The charcoal from two sites has been re-examined by Hazell and Campbell (2019a, 2019b).

Charred grain from three of the site archives held by the museum were examined by Pelling (in prep), and radiocarbon dated in 2017.

## Woodcutts

The Iron Age and Romano-British settlement on Woodcutts Common, associated with tracks and ‘Celtic’ fields, covers about four acres, and was probably a single farmstead. It was excavated by General Pitt Rivers in 1884–5 (Pitt Rivers 1887). Two boxes of charred cereal grain were labelled “grains of corn”, both from pit 43. A total of 536 grain were counted from the two samples. The assemblage was remarkably well preserved and consisted predominantly of *Triticum spelta* (spelt wheat), with a few grains of *Hordeum vulgare* (barley).

Two charred *Triticum spelta* grains were radiocarbon dated from a sample labelled “pit 43, depth 2 feet” (Table 1; Fig. 2). The two radiocarbon determinations are statistically consistent at the 5% significance level ( $T'=2.8$ ;  $v=1$ ;  $T'(5\%)=3.8$ ; Ward and Wilson 1978) and could therefore be of the same age.

## Iwerne

The Iron Age settlement and Roman villa near Park House Farm Buildings, were excavated in 1897 by General Pitt Rivers, but was not published before his death. Much of the archive was discarded or lost, and details are largely known from an examination of a limited surviving archive by Christopher Hawkes and Stuart Piggott (Hawkes et al. 1947). The Iron Age settlement is characterised by numerous pits and early during the Roman period the settlement was modified by the digging of ditches and sub-rectangular pits. A

third century building was speculatively identified as a possible aisled barn. A fourth century building of flint rubble walls, with one room lined with painted wall plaster, was occupied from c AD 300–360. There is no mention of grain in the publication (ibid). One sample of charred grain in the Salisbury Museum collection had lost its original label but was at some point speculatively recorded as from Iwerne. The box contained 2,348 grain, most of which was *Triticum spelta* (spelt wheat) with occasional *Hordeum vulgare* (barley) and occasional possible *Triticum dicoccum* (emmer) and *Bromus* sp. (brome grass).

Three unprovenanced *Triticum spelta* grains were submitted for radiocarbon dating, one of which failed to produce enough carbon for analysis. The two measurements on the grains that were dated (Table 1; Fig. 2) are statistically inconsistent at the 5% significance level ( $T'=10.3$ ;  $v=1$ ;  $T'(5\%)=3.8$ ; Ward and Wilson 1978) and clearly of different ages. The grains are too early to originate from the Roman villa.

## Rotherley

The settlement at Rotherley occupies the brow of a hill and prior to excavation was visible as an irregularly circular enclosure formed by a bank and ditch with an entrance opening into an oblong enclosure on the south-east and associated with further banks and ditches and a field system. Excavations by Pitt Rivers from 1885–6, revealed an Iron Age/Romano-British complex of storage pits with one- or two-house foundations, a corn-drying furnace, foundations of a small rectangular building, granaries and working hollows (Pitt Rivers 1888). Seven samples of charred grain were collected from five features or feature groups: pits 43, 55 and 56, a square hut foundation, and from a group of post holes interpreted as a small raised granary (this last sample is presumably composed of mixed material from more than one post hole). Over 3,500 grain were present in total, dominated by *Hordeum vulgare* (barley) with occasional grain of *Triticum spelta* (spelt wheat), possible *Triticum dicoccum* (emmer wheat), *Bromus* sp. (brome grass) and *Avena* (oats).

Four charred cereal grains were radiocarbon dated from Pit 43 (Table 1; Fig. 2), part of the main circle, and Pit 56, labelled “9 foot 7 inches deep”. The two measurements on *Hordeum* grains from Pit 43 are statistically consistent at the 5% significance level ( $T'=1.6$ ;  $v=1$ ;  $T'(5\%)=3.8$ ; Ward and Wilson 1978) and could therefore be of the same age, while the two measurements on *Hordeum* grains from Pit 56 are statistically inconsistent at the 5% significance level ( $T'=92.9$ ;  $v=1$ ;  $T'(5\%)=3.8$ ; Ward and Wilson 1978) and clearly of different ages.




Figure 2: Probability distribution of the dates on carbonised cereal grains from the Pitt Rivers archive (Woodcutts, Iwerne and Rotherley). The distributions are the result of simple radiocarbon calibration (Stuiver and Reimer 1993)

Table 1: Pitt Rivers archive: Woodcutts, Iwerne, and Rotherley – radiocarbon and stable isotope results

| Laboratory number | Sample reference, material, and context                                                      | $\delta^{13}\text{C}_{\text{IRMS}}$ (‰) | Radiocarbon Age (BP) | Calibrated Date (95% probability) |
|-------------------|----------------------------------------------------------------------------------------------|-----------------------------------------|----------------------|-----------------------------------|
| <b>Woodcutts</b>  |                                                                                              |                                         |                      |                                   |
| UBA-34035         | S.WCT.10.1: sample 1<br>Carbonised grain, <i>Triticum spelta</i> , from Pit 43, depth 2 feet | $-23.6 \pm 0.22$                        | $1855 \pm 33$        | cal AD 90–300                     |
| OxA-35465         | S.WCT.10.1: sample 2<br>Carbonised grain, <i>Triticum spelta</i> , from Pit 43, depth 2 feet | $-24.6 \pm 0.2$                         | $1923 \pm 24$        | cal AD 30–205                     |
| <b>Iwerne</b>     |                                                                                              |                                         |                      |                                   |
| GU43393           | S.WCT.10.1: sample 3<br>Carbonised grain, <i>Triticum spelta</i> , unprovenanced             | Failed: insufficient carbon             |                      |                                   |
| SUERC-73278       | S.WCT.10.1: sample 3b<br>Carbonised grain, <i>Triticum spelta</i> , unprovenanced            | $-23.2 \pm 0.2$                         | $2225 \pm 25$        | 380–200 cal BC                    |
| UBA-34034         | S.WCT.10.1: sample 4<br>Carbonised grain, <i>Triticum spelta</i> , unprovenanced             | $-22.3 \pm 0.22$                        | $2062 \pm 44$        | 200 cal BC–cal AD 60              |

| Laboratory number | Sample reference, material, and context                                                              | $\delta^{13}\text{C}_{\text{IRMS}}$ (‰) | Radiocarbon Age (BP) | Calibrated Date (95% probability) |
|-------------------|------------------------------------------------------------------------------------------------------|-----------------------------------------|----------------------|-----------------------------------|
| <b>Rotherley</b>  |                                                                                                      |                                         |                      |                                   |
| OxA-35464         | S.WCT.10.1: sample 5<br>Carbonised grain, <i>Hordeum</i> ,<br>from the bottom of Pit 43, main circle | -24.1±0.2                               | 2082±24              | 170–5 cal BC                      |
| SUERC-72523       | S.WCT.10.1: sample 6<br>Carbonised grain, <i>Hordeum</i> ,<br>from the bottom of Pit 43, main circle | -24.6±0.2                               | 2034±29              | 135 cal BC–<br>cal AD 60          |
| UBA-34036         | S.WCT.10.1: sample 7<br>Carbonised grain, <i>Hordeum</i> ,<br>from Pit 56, 9 foot 7 inches deep      | -22.9±0.22                              | 1644±35              | cal AD 270–<br>540                |
| OxA-35463         | S.WCT.10.1: sample 8<br>Carbonised grain, <i>Hordeum</i> ,<br>from Pit 56, 9 foot 7 inches deep      | -23.0±0.2                               | 2057±34              | 150 cal BC–<br>cal AD 15          |

## Neolithic and Bronze Age cereals in the Stonehenge landscape

During the review of published archaeobotanical remains from central and southern England (Carruthers and Hunter Dowse 2019; Pelling and Campbell 2013), it became apparent that contamination (intrusion and residuality) was a notable, recurring theme in many assemblages (Pelling et al. 2015). This problem appears to be most prevalent in periods in which plant assemblages are generally less abundant, particularly the Neolithic and early medieval periods. In addition, it was suspected that some assemblages which had not been directly dated, had been attributed to the wrong phase. The importance of directly dating cereal remains from Late Neolithic and Beaker contexts was highlighted in a review of Neolithic and Bronze Age radiocarbon dates by Stevens and Fuller (2012). In order to investigate the degree of intrusion in Neolithic contexts in southern England, and to address insecure phasing of grain assemblages from the Stonehenge landscape, a series of charred grains were dated from three sites in Wiltshire: Durrington Walls, the Cuckoo Stone, and Coneybury Henge.

### Durrington Walls

Durrington Walls is Britain's largest henge enclosure (17ha), lying 3km northeast of Stonehenge in a dry valley perched above the west bank of the River Avon. Previous

excavations (Farrer 1918; Stone et al. 1954; Wainwright and Longworth 1971) demonstrated the considerable quantities of animal bones, Grooved Ware pottery and other occupation debris at this large site, within half an hour's walk of Stonehenge. Excavations in 2004–2007 at the site Walls by the Stonehenge Riverside Project revealed the remains of a large late Neolithic settlement, much of it protected beneath the banks of the henge enclosure (Parker Pearson et al. forth; Parker Pearson 2007; Parker Pearson et al. 2007). An intensive flotation sampling programme produced a modest assemblage of charred plant remains including both wild food plants (*Corylus avellana* (hazel) nutshell, *Malus sylvestris* (crab apple)) and cereal remains (*Triticum dicoccum* grain and glume bases, *Hordeum vulgare* grain) (Simmons forth).

Nine charred cereal grains were dated from three contexts excavated as part of the Stonehenge Riverside Project (Table 2; Fig. 3). All three contexts contain grains of different ages and all are clearly intrusive given the main floruit of activity at the site took place in the middle of the third millennium cal BC (Pelling et al. 2015).




Figure 3: Probability distribution of the dates on carbonised cereal grains from Durrington Walls. The distributions are the result of simple radiocarbon calibration (Stuiver and Reimer 1993)

Table 2: Durrington Walls radiocarbon and stable isotope results

| Laboratory number | Sample, material, and context                                                                      | $\delta^{13}\text{C}_{\text{IRMS}}$ (‰) | Radiocarbon Age (BP) | Calibrated Date (95% probability) |
|-------------------|----------------------------------------------------------------------------------------------------|-----------------------------------------|----------------------|-----------------------------------|
| OxA-27940         | [549] <10079> sample A<br>Carbonised grain, <i>Triticum</i> sp.<br>(free threshing) from [549] the | $-21.7 \pm 0.2$                         | $466 \pm 23$         | cal AD 1419–1456                  |

| Laboratory number | Sample, material, and context                                                                                                                                                                              | $\delta^{13}\text{C}_{\text{IRMS}}$ (‰) | Radiocarbon Age (BP) | Calibrated Date (95% probability) |
|-------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------|----------------------|-----------------------------------|
|                   | fill of a very shallow pit [538]<br>part of the eastern pit complex.                                                                                                                                       |                                         |                      |                                   |
| SUERC-45872       | [549] <10079> sample B<br>Carbonised grain, <i>Triticum</i> sp.<br>(free threshing) from [549] the<br>fill of a very shallow pit [538]<br>part of the eastern pit complex.                                 | $-22.1 \pm 0.2$                         | $169 \pm 34$         | cal AD 1650–<br>1950              |
| OxA-27941         | [585] <10731> sample A<br>Carbonised grain, <i>Hordeum</i><br>indet. (hulled) from [585] a<br>buried soil that formed over the<br>Avenue.                                                                  | $-23.9 \pm 0.2$                         | $1558 \pm 24$        | cal AD 430–<br>575                |
| SUERC-45873       | [585] <10731> sample B<br>Carbonised grain, <i>Hordeum</i><br>indet. (hulled) from [585] a<br>buried soil that formed over the<br>Avenue.                                                                  | $-26.1 \pm 0.2$                         | $1045 \pm 34$        | cal AD 900–<br>1110               |
| UBA-22719         | [585] <10730> sample C<br>Carbonised grain, <i>Hordeum</i><br>indet. from [585] a buried soil<br>that formed over the Avenue.                                                                              | $-24.6 \pm 0.22$                        | $3197 \pm 36$        | 1530–1400<br>cal BC               |
| UBA-22720         | [585] <10730> sample D<br>Carbonised grain, <i>Triticum</i><br><i>dicoccum</i> from [585] a buried<br>soil that formed over the<br>Avenue.                                                                 | $-23.8 \pm 0.22$                        | $3126 \pm 33$        | 1490–1290<br>cal BC               |
| OxA-27942         | [1394] <12067> sample A<br>Carbonised grain, <i>Hordeum</i><br>sp. (hulled) from midden<br>deposit [1394] which<br>accumulated over the<br>abandoned House 902 and<br>was sealed beneath the henge<br>bank | $-22.6 \pm 0.2$                         | $1560 \pm 24$        | cal AD 430–<br>575                |
| SUERC-45874       | [1394] <12067> sample B<br>Carbonised grain, <i>Hordeum</i><br>sp. (hulled) from midden<br>deposit [1394] which<br>accumulated over the<br>abandoned House 902 and<br>was sealed beneath the henge<br>bank | $-23.4 \pm 0.2$                         | $2949 \pm 34$        | 1270–1030<br>cal BC               |
| UBA-22721         | [1394] <12073> sample C<br>Carbonised grain, <i>Triticum</i> sp.                                                                                                                                           | $-23.6 \pm 0.22$                        | $78 \pm 33$          | cal AD 1680–<br>1930              |

| Laboratory number | Sample, material, and context                                                                                                    | $\delta^{13}\text{C}_{\text{IRMS}}$ (‰) | Radiocarbon Age (BP) | Calibrated Date (95% probability) |
|-------------------|----------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------|----------------------|-----------------------------------|
|                   | (free threshing) from midden deposit [1394] which accumulated over the abandoned House 902 and was sealed beneath the henge bank |                                         |                      |                                   |

## Cuckoo Stone

The recumbent sarsen stone known as the Cuckoo Stone lies 2.6km northeast of Stonehenge. Two pits, Pit 135 and Pit 180, were excavated as part of the Stonehenge Riverside Project (Parker Pearson et al. 2020). Radiocarbon dates on a cattle scapula shovel (OxA-18940; 4253±28 BP; Parker Pearson et al. 2020, table 7.2) from Pit 135 and articulating red deer bones (SUERC-46473; 4231±27 BP; Parker Pearson et al. 2020, table 7.2) from Pit 180 clearly demonstrate they were dug and infilled in the first quarter of the third millennium and are thus Late Neolithic.

Eleven cereal grains and one glume wheat glume base were recovered from fill (181) of Pit 180 and 11 cereal grains and one probable free-threshing, wheat rachis node from fill (136) of Pit 135 (Simmons 2020). In order to determine whether the cereals were Late Neolithic in date, or intrusive, two grains (*Hordeum* sp. and free-threshing *Triticum* sp.) were dated from each pit (Table 3; Fig. 4). The results clearly indicate that the grains are intrusive and represent material that has ‘contaminated’ the pit at different times (Simmons 2020, 390).




Figure 4: Probability distribution of the dates on carbonised cereal grains from the Cuckoo Stone. The distributions are the result of simple radiocarbon calibration (Stuiver and Reimer 1993)

Table 3: Cuckoo Stone radiocarbon and stable isotope measurements

| Laboratory number | Sample, material, and context                                                                                            | Radiocarbon age (BP) | $\delta^{13}\text{C}_{\text{AMS}}$ (‰) | $\delta^{13}\text{C}_{\text{IRMS}}$ (‰) | Calibrated Date (95% probability) |
|-------------------|--------------------------------------------------------------------------------------------------------------------------|----------------------|----------------------------------------|-----------------------------------------|-----------------------------------|
| ETH-89715         | 102. <i>Hordeum</i><br>Carbonised grain, <i>Hordeum</i> sp. indet. from the fill (136) of Pit 135                        | 1681±23              | -23.0                                  |                                         | cal AD 260–420                    |
| GrM-14843         | 102. <i>Triticum</i><br>Carbonised grain, cf. <i>Triticum</i> sp. (free threshing wheat) from the fill (136) of Pit 135  | 670±20               |                                        | -23.0±0.15                              | cal AD 1283–1386                  |
| ETH-89714         | 114. <i>Hordeum</i><br>Carbonised grain, <i>Hordeum</i> sp. indet. from the fill (180) of Pit 181                        | 1732±22              | -21.5                                  |                                         | cal AD 250–400                    |
| GrM-14841         | 115. <i>Triticum</i><br>Carbonised grain, cf. <i>Triticum</i> sp. (free threshing wheat), from the fill (180) of Pit 181 | 810±20               |                                        | -23.0±0.15                              | cal AD 1209–1269                  |

## Coneybury Henge

A henge, surviving as a cropmark and initially recorded as a ploughed-out disc barrow Coneybury Hill, Coneybury Henge was subject to surface collection, geophysical survey and excavation in 1980 as part of the Stonehenge Environs Project (Richards 1990). The whole site slopes gently to the south, and the interior of the enclosure appears to have been cut back into the hillside in order to create a level interior platform. Excavation confirmed that the enclosing ditch, with a single entrance to the north east, had been accompanied by an external bank. Various internal features were excavated, comprising a few pits and post holes, numerous stake holes, and an arc of post holes concentric to the inner edge of the enclosure ditch which may have represented a post-circle. Some of the internal features probably pre-dated enclosure construction, and pottery from the site suggests that activity spanned the Early Neolithic through to the Middle Bronze Age. At the entrance, the one ditch terminal to be excavated contained a large, apparently cumulative deposit, including a substantial quantity of cattle bones and lithic material indicative of carcass preparation and cooking associated with Beaker pottery.

Two sections were excavated across the ditch, one to the south and the other across the western terminal. Radiocarbon dating of the part skeleton of a male dog from the primary

fill of the ditch (OxA-1408;  $4200 \pm 110$  BP) suggests it was dug in the first half of the second millennium cal BC.

The fill of the ditch contained a sample of just over 300 naked and hulled barley grains in the ratio of 3:2 (Carruthers 1990). While the specialist report suggested the grain deposit was recovered from a secondary, Early Bronze Age deposit (Carruthers 1990, 250), it is given as Late Neolithic in the table (Carruthers 1990, 251). A selection of three naked and three hulled barley grains were dated to confirm the age of the deposit. The six radiocarbon determinations on individual barley grains (Table 4) are statistically consistent at the 5% significance level ( $T'=6.3$ ;  $v=5$ ;  $T'(5\%)=11.1$ ; Ward and Wilson 1978) and could therefore be of the same actual age. The barley found in the ditch therefore dates to the centuries either side of 2000 cal BC and is hence much later than the primary infilling of the ditch (Fig. 5) and is not related to Late Neolithic activity at Coneybury Henge (Pelling and Campbell 2013).




Figure 5: Probability distribution of the dates on carbonised cereal grains from Coneybury Henge. The distributions are the result of simple radiocarbon calibration (Stuiver and Reimer 1993)

Table 4: Coneybury Henge radiocarbon and stable isotope measurements

| Laboratory number | Sample, material, and context                                                                                                                    | $\delta^{13}\text{C}_{\text{IRMS}}$ (‰) | Radiocarbon Age (BP) | Calibrated Date (95% probability) |
|-------------------|--------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------|----------------------|-----------------------------------|
| UBA-22185         | 1444 (naked) sample A<br>Carbonised grain; <i>Hordeum vulgare</i> var <i>nudeum</i> (naked six-row barley) from the fill of the henge ditch (W2) | $-24.4 \pm 0.22$                        | $3594 \pm 37$        | 2110–1800 cal BC                  |

| Laboratory number | Sample, material, and context                                                                                                                    | $\delta^{13}\text{C}_{\text{IRMS}}$ (‰) | Radiocarbon Age (BP) | Calibrated Date (95% probability) |
|-------------------|--------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------|----------------------|-----------------------------------|
| SUERC-44490       | 1444 (naked) sample B<br>Carbonised grain; <i>Hordeum vulgare</i> var <i>nudeum</i> (naked six-row barley) from the fill of the henge ditch (W2) | $-24.7 \pm 0.2$                         | $3611 \pm 27$        | 2110–1890 cal BC                  |
| OxA-27614         | 1444 (naked) sample V<br>Carbonised grain; <i>Hordeum vulgare</i> var <i>nudeum</i> (naked six-row barley) from the fill of the henge ditch (W2) | $-24.4 \pm 0.2$                         | $3695 \pm 30$        | 2190–1980 cal BC                  |
| UBA-22184         | 1444 (hulled) sample A<br>Carbonised grain; <i>Hordeum vulgare</i> L emend (hulled six-row barley) from the fill of the henge ditch (W2)         | $-26.3 \pm 0.22$                        | $3640 \pm 33$        | 2130–1900 cal BC                  |
| SUERC-44489       | 1444 (hulled) sample B<br>Carbonised grain; <i>Hordeum vulgare</i> L emend (hulled six-row barley) from the fill of the henge ditch (W2)         | $-24.3 \pm 0.2$                         | $3620 \pm 29$        | 2120–1890 cal BC                  |
| OxA-27707         | 1444 (hulled) sample C<br>Carbonised grain; <i>Hordeum vulgare</i> L emend (hulled six-row barley) from the fill of the henge ditch (W2)         | $-24.5 \pm 0.22$                        | $3639 \pm 29$        | 2140–1920 cal BC                  |

## Contamination: archaeophytes

A review of archaeophytes (plant species introduced during the Holocene before AD 1500) indicated an increasing number of species arrived in the British Isles through time, with a number of classic cornfield weeds including *Anthemis cotula* (stinking mayweed) and *Agrostemma githago* (corncockle) arriving in the late Iron Age or Roman period (Preston et al. 2004). Given the inherent risk of contamination of deposits by more recent material, finds of a particular taxon significantly earlier than usual should be treated with caution. (Pelling et al. 2015).

## Thanet Earth

Situated between Monkton and Birchington (near Margate) on the Isle of Thanet, Kent, the ‘Thanet Earth’ site covers 90 hectares (about 220 acres), of which about 47 hectares (nearly half a million square metres) have been examined archaeologically. Open area excavations in 2007–8 prior to the building of the UK’s biggest glasshouse complex by the

Canterbury Archaeological Trust revealed nine barrows, 63 structures, 75 sunken featured structures, about 70 enclosures and 33 trackways (including drove ways and hollow ways), and a considerable number of burials (both inhumation and cremation burials, all of prehistoric or Roman provenance) (Rady 2010).

The fill of pit [10454] contained a surprisingly high concentration of grains of *Triticum aestivum/turgidum* (free-threshing wheat), one of which was directly dated to the Early Neolithic (UBA-22207; 5000±45 BP), as well as tetraploid wheat rachis identified as a variety of *Triticum turgidum* (rivet wheat type) (Carruthers 2019; Henshaw et al. 2019, table 6). Also present in the pit were *Corylus avellana* (hazel) nutshell fragments, seeds of *Anthemis cotula* (stinking chamomile), a possible capsule fragment from corncockle (*Agrostemma githago*) and a possible *Pisum sativum* (pea). *Anthemis cotula* and *Agrostemma githago* are considered archaeophytes, introduced to the British Isles in the Late Iron Age or Roman period (Preston et al. 2004, table 2), while pulses are not reliably recorded prior to the Middle Bronze Age (Treasure and Church 2016). The radiocarbon dating of *Anthemis cotula* seeds (Table 5) was undertaken to investigate the extent of contamination of the pits fill.

The fact that the *Anthemis cotula* seeds dated to the first two centuries of the second millennium cal AD (Fig. 5) confirms that they are intrusive despite the pit being apparently well sealed and containing Early Neolithic grain.




Figure 6: Probability distribution of the date on plant macrofossils from Thanet Earth. The distribution is the result of simple radiocarbon calibration (Stuiver and Reimer 1993)

Table 5: Thanet Earth radiocarbon and stable isotope measurements

| Laboratory number | Sample, material & context                                                                                           | $\delta^{13}\text{C}$ (‰) | Radiocarbon Age (BP) | Calibrated Date (95% probability) |
|-------------------|----------------------------------------------------------------------------------------------------------------------|---------------------------|----------------------|-----------------------------------|
| UBA-25299         | Context (10452) <1851><br>Carbonised plant macrofossil, <i>Anthemis cotula</i> x 9, from fill (10452) of pit [10454] | –                         | 948±38               | cal AD 1020–1200                  |

## Regional Reviews of charred plant remains

English Heritage (now Historic England) has commissioned three regional reviews of plant macrofossil data covering the northern, midland, and southern counties of England. The first of these, '*A Review of the Evidence for Macrofossil Plant Remains from Archaeological Deposits in Northern England*', was produced by Hall and Huntley (2007). In support of the midlands regional review (Carruthers and Hunter Dowse 2019) radiocarbon dating was undertaken on material from Lockington, Leicestershire.

### Lockington

Excavations in 1994 in advance of the construction of the A564(T) Derby southern bypass near the village of Lockington in north-west Leicestershire exposed an early Bronze Age barrow cemetery. Prior to the fieldwork a cluster of cropmarked features at Site V suggested the presence of a pit circle (Hughes 2000). The fill (175) of F182 was the most productive, in terms of charred plant remains, of the features excavated from Site V. The presence of *T. dicoccum* (emmer) and *Triticum spelta* (spelt) were confirmed from diagnostic chaff (glume bases) forming 62% of the recovered charred material by count (Moffett and Monkton 2000). Radiocarbon dating of *Corylus avellana* and *Prunus spinosa* charcoal (hazel and blackthorn) from the pit (Beta-83721; 3039±80 BP) suggested it was infilled in the middle Bronze Age (1470–1040 cal BC; 95% probability).

In order to confirm the potential early occurrence of spelt wheat in the region four *Triticum spelta* glume bases from pit F182 were dated (Table 6). The glume bases are noticeably younger than the charcoal (Fig 7) and given the high proportion of chaff in the feature, suggesting that it is the by-product from the preparation of wheat grain (Hillman 1981), it is probable that the charcoal is residual. While the spelt wheat was the earliest occurrence of spelt wheat in Leicestershire at the time of dating, it is significantly later than the introduction of spelt to the southern counties where it is recorded from the late Early/Middle Bronze Age (Carruthers and Hunter Dowse, 2019, 199) and is unlikely to be the earliest example in the midlands.




Figure 7: Probability distribution of the dates from Lockington F182. The distributions are the result of simple radiocarbon calibration (Stuiver and Reimer 1993)

Table 6: Lockington radiocarbon and stable isotope measurements

| Laboratory number | Sample, material and context                                                                                       | $\delta^{13}\text{C}$ (‰) | Radiocarbon Age (BP) | Calibrated Date (95% probability) |
|-------------------|--------------------------------------------------------------------------------------------------------------------|---------------------------|----------------------|-----------------------------------|
| UBA-25298         | Context [175] <38><br>Carbonised plant macrofossil, <i>Triticum spelta</i> glume base x 4, from fill (175) of F182 | –                         | 2737 $\pm$ 38        | 990–810 cal BC                    |

## Hunsbury

The monument at Hunsbury is situated on the summit of a prominent hill overlooking the Nene Valley, Northamptonshire, and includes the earthwork and buried remains of a multivallate Iron Age hillfort (Fell 1936). Finds recovered during excavations for iron stone at the end of the 19th century are held by the Northants and British museums. A box of charred cereal grain recovered from the site had been held by the Economic Botany collection at Kew since it was deposited by T J George of the Northampton Public Museum in 1902. The box label states that the grain was collected from a quern. The box of grain was forwarded to Historic England for scientific analysis (Bouchard-Perron forthcoming).

In order to confirm the assumed middle Iron Age date of the charred grain deposit from Hunsbury three grains of *Triticum spelta* were dated (Table 7). The radiocarbon determinations are statistically consistent at the 5% significance level ( $T'=1.3$ ,  $T'(5\%)=3.8$ ,  $v=2$ ; Ward and Wilson 1978) and could be of the same actual age. The results confirm the grains are middle Iron Age in date (Fig. 8).




Figure 8: Probability distribution of the dates on carbonised cereal grains from Hunsbury hillfort. The distributions are the result of simple radiocarbon calibration (Stuiver and Reimer 1993)

Table 7: Hunsbury radiocarbon and stable isotope measurements

| Laboratory Code | Sample, material & depth | $\delta^{13}\text{C}_{\text{IRMS}}$ (‰) | $\delta^{13}\text{C}_{\text{AMS}}$ (‰) | Radiocarbon Age (BP) | Calibrated Date (95% probability) |
|-----------------|--------------------------|-----------------------------------------|----------------------------------------|----------------------|-----------------------------------|
|-----------------|--------------------------|-----------------------------------------|----------------------------------------|----------------------|-----------------------------------|

|            |                                                                                                                       |            |           |         |                   |
|------------|-----------------------------------------------------------------------------------------------------------------------|------------|-----------|---------|-------------------|
| GrM-19929  | 26786.1<br>Carbonised grain,<br><i>Triticum spelta</i> , from<br>Kew Economic Botany<br>collection (Cat. N.<br>26786) | -23.5±0.15 |           | 2289±24 | 390–210 cal<br>BC |
| ETH-102412 | 26786.2<br>Carbonised grain,<br><i>Triticum spelta</i> , from<br>Kew Economic Botany<br>collection (Cat. N.<br>26786) |            | -23.3±1.0 | 2252±22 | 390–210 cal<br>BC |
| ETH-102413 | 26786.3<br>Carbonised grain,<br><i>Triticum spelta</i> , from<br>Kew Economic Botany<br>collection (Cat. N.<br>26786) | -25.3±1.0  |           | 2263±22 | 395–205 cal<br>BC |

## References

- Aerts-Bijma, A. T., Paul, D., Dee, M. W., Palstra, S. W. L., and Meijer, H. A. J. 2021 'An independent assessment of uncertainty for radiocarbon analysis with the new generation high-yield accelerator mass spectrometers', *Radiocarbon*, 63(1), 1–22: <https://doi.org/10.1017/RDC.2020.101>
- Bouchard-Perron, J-A. forthcoming Hunsbury Hill, Northamptonshire: Analysis of 19th century excavation plant remains', Historic England Research Report Series.
- Brock, F., Higham, T., Ditchfield, P., and Bronk Ramsey, C. 2010 'Current pretreatment methods for AMS radiocarbon dating at the Oxford Radiocarbon Accelerator Unit (ORAU)', *Radiocarbon*, 52, 103–12: <https://doi.org/10.1017/S0033822200045069>
- Bronk Ramsey, C. 1995 'Radiocarbon calibration and analysis of stratigraphy: the OxCal program', *Radiocarbon*, 37(2), 425–30: <https://doi.org/10.1017/S0033822200030903>
- Bronk Ramsey, C. 2001 'Development of the radiocarbon calibration program', *Radiocarbon*, 43, 355–63: <https://doi.org/10.1017/S0033822200038212>
- Bronk Ramsey, C. 2009 'Bayesian analysis of radiocarbon dates', *Radiocarbon*, 51(1), 337–60: <https://doi.org/10.1017/S0033822200033865>
- Bronk Ramsey, C., Ditchfield, P., and Humm, M. 2004 'AMS methods and developments - using a gas ion source for radiocarbon AMS and GC-AMS', *Radiocarbon*, 46, 25–33: <https://doi.org/10.1017/S003382220003931X>
- Carruthers, W. 1990 'Carbonised plant remains' in Richards, J. ed. *The Stonehenge Environs Project*, Archaeological Report (London), 250–2.
- Carruthers, W. J. 2019 'Charred, waterlogged and mineralised plant remains', in Henshaw, R., Holman, J., Masfield, R., Rady, J. and Weekes, J. eds *Beneath the Seamark: 6000 years of an Island's History. Archaeological Investigations at Thanet Earth, Kent 2007-2012, Volume 2: Specialist Reports*. Canterbury Archaeological Trust Technical Report 2 (Canterbury), 1087–1135.
- Carruthers, W. J., and Hunter Dowse, K. L. 2019 'A Review of Macroscopic Plant Remains from the Midland Counties', Historic England Research Report Series 47-2019: <https://historicengland.org.uk/research/results/reports/47-2019> (acc. 15 May 2023).

- Dee, M., and Bronk Ramsey, C. 2000 'Refinement of graphite target production at ORAU', *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms*, 172, 449–53: [https://doi.org/10.1016/S0168-583X\(00\)00337-2](https://doi.org/10.1016/S0168-583X(00)00337-2)
- Dee, M. W., Palstra, S. W. L., Aerts-Bijma, A. T., Bleeker, M. O., De Bruijn, S., Ghebru, F., Jansen, H. G., Kuitens, M., Paul, D., Richie, R. R., Spriensma, J. J., Scifo, A., Van Zonneveld, D., Verstappen-Dumoulin, B. M. a. A., Wietzes-Land, P., and Meijer, H. a. J. 2019 'Radiocarbon dating at Groningen: New and updated chemical pretreatment procedures', *Radiocarbon*, 62(1), 63–74: <https://doi.org/10.1017/RDC.2019.101>
- Dunbar, E., Cook, G. T., Naysmith, P., Tipney, B. G., and Xu, S. 2016 'AMS <sup>14</sup>C dating at the Scottish Universities Environmental Research Centre (SUERC) Radiocarbon Dating Laboratory', *Radiocarbon*, 58, 9–23: <https://doi.org/10.1017/RDC.2015.2>
- Farrer, P. 1918 'Durrington Walls or Long Walls', *Wiltshire Archaeological and Natural History Magazine*, 40, 95–103.
- Fell, C. I. 1936 'The Hunsbury hill-fort, Northants: a new survey of the material', *Archaeological Journal*, 93, 57–100: <https://doi.org/10.1080/00665983.1936.10853661>
- Hajdas, I. 2008 'Radiocarbon dating and its applications in Quaternary studies', *Eiszeitalter und Gegenwart Quaternary Science Journal*, 57, 2–24: <https://doi.org/10.3285/eg.57.1-2.1>
- Hall, A. R., and Huntley, J. P. 2007 'A review of the evidence for macrofossil plant remains from archaeological deposits in Northern England', Historic England Research Report Series 87-2007: <https://historicengland.org.uk/research/results/reports/87-2007> (acc. 15 May 2023).
- Hazell, Z., and Campbell, G. 2019a 'Woodcutts Common, Cranborne Chase, Dorset. Analysis of wood charcoal remains from the Pitt Rivers Archive, The Salisbury Museum' Historic England Research Report Series 57-2018: <https://historicengland.org.uk/research/results/reports/57-2018> (acc. 15 May 2023)
- Hazell, Z., and Campbell, G. 2019b 'Rotherley, Cranborne Chase, Wiltshire. Analysis of wood charcoal remains from the Pitt Rivers Archive', The Salisbury Museum Research Historic England Research Report Series 59-2018: <https://historicengland.org.uk/research/results/reports/58-2018> (acc. 15 May 2023)
- Hawkes, C. F. C., Piggott, S., and Gray, H. St. G. 1947 'Britons, Romans and Saxon around Salisbury and in Cranborne Chase part II, reviewing the excavations of General Pitt Rivers, 1881–1897', *The Archaeological Journal*, 104, 27–81

- Henshaw, R., Holman, J., Masefield, R., Rady, J., and Weekes, J. eds 2019 *Beneath the Seamark: 6000 years of an Island's History. Archaeological Investigations at Thanet Earth, Kent 2007–2012, Volume 5: Tables*, Canterbury Archaeological Trust Technical Report 2 (Canterbury).
- Hillman, G. 1981 'Reconstructing crop husbandry practices from charred remains of crops, in' Mercer, R. ed *Farming Practice in British Prehistory* (Edinburgh), 123–62.
- Hughes, G. 2000 *The Lockington Gold Hoard: an early Bronze Age cemetery at Lockington, Leicestershire* (Oxford).
- Moffett, L., and Monckton, A. 2000 'Charred plant remains', in Hughes, G. ed. *The Lockington Gold Hoard: An Early Bronze Age Barrow Cemetery at Lockington, Leicestershire* (Oxford), 78–81.
- Mook, W. G. 1986 'Business meeting: Recommendations/Resolutions adopted by the Twelfth International Radiocarbon Conference', *Radiocarbon*, 28, 799:  
<https://doi.org/10.1017/S0033822200008043>
- Parker Pearson, M. 2007 'The Stonehenge Riverside Project: excavations at the east entrance of Durrington Walls', in Larsson, M and Parker Pearson, M. eds. *From Stonehenge to the Baltic*, British Archaeological Reports International Series 1692 (Oxford) 125–44.
- Parker Pearson M., Cleal R., Marshall, P., Needham, S., Pollard, J., Richards, C., Ruggles, C., Sheridan, A., Thomas, J., Tilley, C., Welham, K., Chamberlain, A., Chenery, C., Evans, J., Knüsel C., Linford, N., Martin, L., Montgomery, J., Payne, A., and Richards, M. 2007 'The age of Stonehenge', *Antiquity*, 81, 617–39:  
<https://doi.org/10.1017/S0003598X00095624>
- Parker Pearson, M., Pollard, J., Richards, C., Thomas, J., Tilley, C., and Welham, K. 2020. *Stonehenge for the Ancestors. Part 1: Landscape and Monument* (Leiden).
- Parker Pearson, M., Pollard, J., Richards, C., Thomas, J., Tilley, C., and Welham, K. forthcoming *Durrington Walls and Woodhenge: A place for the living, The Stonehenge Riverside Project Volume 3* (Leiden).
- Pelling, R., and Campbell C. 2013 'Plant resources' in Canti, M., Campbell, G., and Greaney, S. eds 'Stonehenge World Heritage Site Synthesis: Prehistoric Landscape, Environment and Economy' Historic England Research Report Series 45-2013:  
<https://historicengland.org.uk/research/results/reports/45-2013> (acc. 15 May 2023) 37–60.

Pelling, R., Campbell, G., Carruthers, W., Hunter, K., and Marshall, P. 2015 'Exploring contamination (intrusion and residuality) in the archaeobotanical record: case studies from central and southern England', *Vegetation History and Archaeobotany*, 24, 85–99: <https://doi.org/10.1007/s00334-014-0493-8>

Pitt Rivers, A. 1887 *Excavations in Cranborne Chase near Rushmore, on the borders of Dorset and Wilts. Vol. I: excavations in the Romano-British village on Woodcutts Common, and Romano-British antiquities in Rushmore Park.*

Pitt Rivers, A. 1888 *Excavations in Cranborne Chase, near Rushmore, on the borders of Dorset and Wilts 1880–1888. Vol. II: Excavations in Barrows near Rushmore. Excavations in Romano-British Village, Rotherley. Excavations in Winkelbury Camp. Excavations in British Barrows and Anglo-Saxon Cemetery, Winkelbury Hill.*

Preston, C. D., Pearman, D. A., and Hall, A. R. 2004 'Archaeophytes in Britain', *Botanical Journal of the Linnean Society*, 145(3), 257–94: <https://doi.org/10.1111/j.1095-8339.2004.00284.x>

Rady, J. 2010 'Excavations at Thanet Earth 2007–2008: Assessment Report' (Unpublished report, Canterbury Archaeological Trust).

Reimer, P. J., Austin, W. E. N., Bard, E., Bayliss, A., Blackwell, P. G., Bronk Ramsey, C., Butzin, M., Cheng, H., Edwards, R. L., Friedrich, M., Grootes, P. M., Guilderson, T. P., Hajdas, I., Heaton, T. J., Hogg, A. G., Hughen, K. A., Kromer, B., Manning, S. W., Muscheler, R., Palmer, J. G., Pearson, C., Van Der Plicht, J., Reimer, R. W., Richards, D. A., Scott, E. M., Southon, J. R., Turney, C. S. M., Wacker, L., Adolphi, F., Büntgen, U., Capano, M., Fahrni, S. M., Fogtmann-Schulz, A., Friedrich, R., Köhler, P., Kudsk, S., Miyake, F., Olsen, J., Reinig, F., Sakamoto, M., Sookdeo, A., and Talamo, S. 2020 'The IntCal20 northern hemisphere radiocarbon age calibration curve (0–55 cal kBP)', *Radiocarbon*, 64(2), 725–57: <https://doi.org/10.1017/RDC.2020.41>

Reimer, P. J., Hoper, S., McDonald, J., Reimer, R., Svyatko, S. and Thompson, M. 2015. 'The Queen's University, Belfast: Laboratory Protocols used for AMS Radiocarbon Dating at the <sup>14</sup>CHRONO Centre', Historic England Research Report Series 5-2015: <https://historicengland.org.uk/research/results/reports/5-2015> (acc. 15 May 2023).

Richards, J. 1990 *The Stonehenge Environs Project* (London).

Salehpour, M., Håkansson, K., Possnert, G., Wacker, L., and Synal, H.-A. 2016 'Performance report for the low energy compact accelerator mass spectrometer at Uppsala University', *Nuclear Instruments and Methods in Physics Research Section B: Beam*

*Interactions with Materials and Atoms*, 371, 360–4:

<https://doi.org/10.1016/j.nimb.2015.10.034>

Scott, E. M., Naysmith, P., and Cook, G. T. 2017 'Should Archaeologists Care about  $^{14}\text{C}$  Intercomparisons? Why? A Summary Report on SIRI', *Radiocarbon*, 59(5), 1589–96:

<https://doi.org/10.1017/RDC.2017.12>

Simmons, E. 2020 'Charred plant remains from around the Cuckoo Stone' in Parker Pearson, M., Pollard, J., Richards, C., Thomas, J., Tilley, C., and Welham, K. eds *Stonehenge for the Ancestors: Landscape and Monuments, The Stonehenge Riverside Project Volume 1* (Leiden) 390–91.

Simmons, E. forthcoming 'Plant remains and wood charcoal' in Parker Pearson, M., Pollard, J., Richards, C., Thomas, J., Tilley, C., and Welham, K. eds *Durrington Walls and Woodhenge: A place for the living, The Stonehenge Riverside Project Volume 3* (Leiden).

Sookdeo, A., Kromer, B., Büntgen, U., Friedrich, M., Friedrich, R., Helle, G., Pauly, M., Nievergelt, D., Reinig, F., Treydte, K., Synal, H.-A., and Wacker, L. 2020 'Quality Dating: A well-defined protocol implemented at ETH for high-precision  $^{14}\text{C}$  dates tested on Late Glacial wood', *Radiocarbon*, 62(4), 891–9: 10.1017/RDC.2019.132

Stevens, C. J., and Fuller, D. Q. 2012 'Did Neolithic farming fail? The case for a Bronze Age agricultural revolution in the British Isles', *Antiquity* 86, 707–22:

<https://doi.org/10.1017/S0003598X00047864>

Stone, J. F. S., Piggott, S., and Booth, A. S. J. 1954 'Durrington walls, Wiltshire: Recent excavations at a ceremonial site of the early second millennium B.C.', *The Antiquaries Journal*, 34, 155–77: <https://doi.org/10.1017/S0003581500059837>

Stuiver, M., and Polach, H. A. 1977 'Discussion reporting of  $^{14}\text{C}$  data', *Radiocarbon*, 19(3), 355–63: <https://doi.org/10.1017/S0033822200003672>

Stuiver, M., and Reimer, P. J. 1993 'Extended  $^{14}\text{C}$  data base and revised CALIB 3.0  $^{14}\text{C}$  age calibration program', *Radiocarbon*, 35, 215–30:

<https://doi.org/10.1017/S0033822200013904>

Synal, H.-A., Stocker, M., and Suter, M. 2007 'MICADAS: A new compact radiocarbon AMS system', *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms*, 259, 7–13:

<https://doi.org/10.1016/j.nimb.2007.01.138>

Treasure, E. R., and Church, M. J. 2016 'Can't find a pulse? Celtic bean (*Vicia faba* L.) in British prehistory', *Environmental Archaeology*, 22(2), 113–27:

<https://doi.org/10.1080/14614103.2016.1153769>

Vogel, J. S., Southon, J. R., Nelson, D. E., and Brown, T. A. 1984 'Performance of catalytically condensed carbon for use in accelerator mass-spectrometry', *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms*, 233, 289–93: [https://doi.org/10.1016/0168-583X\(84\)90529-9](https://doi.org/10.1016/0168-583X(84)90529-9)

Wainwright, G. J., and Longworth, I. H. 1971 *Durrington Walls: Excavations 1966–1968*, Reports of the Research Committee of the Society of Antiquaries of London 29 (London).

Wacker, L., Němec, M., and Bourquin, J. 2010a 'A revolutionary graphitisation system: fully automated, compact and simple', *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms*, 268, 931–4:

<https://doi.org/10.1016/j.nimb.2009.10.067>

Wacker, L., Bonani, G., Friedrich, M., Hajdas, I., Kromer, B., Němec, M., Ruff, M., Suter, M., Synal, H-A., and Vockenhuber, C. 2010b 'MICADAS: routine and high-precision radiocarbon dating', *Radiocarbon*, 52, 252–62:

<https://doi.org/10.1017/S0033822200045288>

Wacker, L., Christl, M., and Synal, H-A. 2010c 'Bats: A new tool for AMS data reduction', *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms*, 268, 976–9: <https://doi.org/10.1016/j.nimb.2009.10.078>

Ward, G. K. and Wilson, S. R. 1978 'Procedures for comparing and combining radiocarbon age determinations: a critique', *Archaeometry*, 20, 19–32. <https://doi.org/10.1111/j.1475-4754.1978.tb00208.x>



Historic England

# Historic England's Research Reports

We are the public body that helps people care for, enjoy and celebrate England's historic environment.

We carry out and fund applied research to support the protection and management of the historic environment. Our research programme is wide-ranging and both national and local in scope, with projects that highlight new discoveries and provide greater understanding, appreciation and enjoyment of our historic places.

More information on our research strategy and agenda is available at

[HistoricEngland.org.uk/research/agenda](https://HistoricEngland.org.uk/research/agenda).

The Research Report Series replaces the former Centre for Archaeology Reports Series, the Archaeological Investigation Report Series, the Architectural Investigation Report Series, and the Research Department Report Series.

All reports are available at [HistoricEngland.org.uk/research/results/reports](https://HistoricEngland.org.uk/research/results/reports). There are over 7,000 reports going back over 50 years. You can find out more about the scope of the Series here: [HistoricEngland.org.uk/research/results/about-the-research-reports-database](https://HistoricEngland.org.uk/research/results/about-the-research-reports-database).

Keep in touch with our research through our digital magazine *Historic England Research*

[HistoricEngland.org.uk/whats-new/research](https://HistoricEngland.org.uk/whats-new/research).